
# 2012 KENTUCKY ASTHMA PROGRAM FACT SHEET: BARREN RIVER AREA DEVELOPMENT DISTRICT


- The Barren River District of Kentucky is comprised of 10 counties: Logan, Simpson, Butler, Warren, Edmonson, Hart, Barren, Allen, Metcalfe, and Monroe.
- The 2011 Barren River District population was roughly 286,258. About 66,700 persons were younger than 18 (children), and about 219,600 were 18 or older (adults).



## **PREVALENCE**

 In 2010, an estimated 6,200 children (9.6%) in the Barren River District currently had asthma, compared to 10.7% for Kentucky and 8.4% for the United States.<sup>1</sup> In 2011, an estimated 23,000 adults (10.8%) in the Barren River District currently had asthma, compared to 10.4% for Kentucky and 9.1% for the United States.<sup>1</sup>







 In 2010, child current asthma prevalence rate was lower in the Barren River District compared to Kentucky. In 2011, adult current asthma prevalence rate was higher among females than males in this district. This pattern was similar for Kentucky. (NOTE: BRFSS 2010 data cannot be compared to BRFSS 2011 data because of change in methodology)

### **HOSPITALIZATIONS**

 In 2011, the age-adjusted hospitalization rate for children was 3.1/10,000 persons compared to the Kentucky rate of 4.9/10,000 persons; while the age-adjusted hospitalization rate for adults was 13.7/10,000 persons compared to the Kentucky rate of 9.7/10,000 persons.





Overall, child asthma hospitalizations have been lower in Barren River District compared to Kentucky while adult
hospitalization rates have been higher in this district compared to Kentucky. Between 2001 and 2011, the Barren River
District averaged 10.0 total asthma hospitalizations/10,000 persons/year, while Kentucky averaged 8.3 total
hospitalizations/10,000 persons/ year over the same 11 year period.<sup>2</sup>

#### **EMERGENCY DEPARTMENT VISITS**

• In 2011, the asthma Emergency Department (ED) visits for children were 1.5/1,000 persons, compared to the state rate of 1.9/1,000 persons<sup>3</sup> and for adults were 1.7/1,000 persons compared to the state rate of 2.5/1,000 persons.<sup>3</sup>

#### **MORTALITY**

Between 2001 and 2010, the average age-adjusted asthma deaths were 0.94 /100,000 persons/year in this district compared to 1.10/100,000 persons/year for Kentucky.<sup>4</sup> In 2010, crude asthma death rates for Kentucky were 1.01/100,000 persons<sup>4</sup> compared to the US crude death rate of 1.1/100,000 persons.<sup>5</sup>

#### NOTES:

- 1. Behavioral Risk Factor Surveillance System (BRFSS).
- 3. OHP: Preliminary data 2011.
- 5. CDC Wonder data: http://wonder.cdc.gov/ucd-icd10.html
- 2. Kentucky, Office of Health Policy (OHP), 2001-2011.
- 4. Office of Vital Statistics, Kentucky, 2001-2010.

This fact sheet was supported by the Cooperative Agreement Number 5U59EH000508-04 from the Centers for Disease Control and Prevention (CDC). Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the CDC.

Jan Hatfield, Program Manager; Sue Thomas-Cox, PI; Deepa Valvi, Asthma Epidemiologist 275 E. Main St. \* Mail Stop HS2WE \* Frankfort, KY 40621 \* (502) 564-7996, ext. 4439

Website: www.chfs.ky.gov/Asthma